

The role of afterglow break–times as GRB jet angle indicators

L. Nava^{1,2*}, G. Ghisellini¹, G. Ghirlanda¹, J.I. Cabrera³, C. Firmani^{1,3} and V. Avila–Reese³

¹Osservatorio Astronomico di Brera, via E. Bianchi 46, I-23807 Merate, Italy

²Università degli Studi dell’Insubria, Dipartimento di Fisica e Matematica, via Valleggio 11, I-22100 Como, Italy

³Instituto de Astronomía, Universidad Nacional Autónoma de México, A.P. 70-264, 04510, México, D.F., México

Accepted 2007 February 27. Received 2007 February 23; in original form 2007 January 24

ABSTRACT

The early X–ray light curve of Gamma Ray Bursts (GRBs) is complex, and shows a typical steep–flat–steep behaviour. The time T_a at which the flat (plateau) part ends may bear some important physical information, especially if it plays the same role of the so called jet break time t_{jet} . To this aim, stimulated by the recent analysis of Willingale et al., we have assembled a sample of GRBs of known redshifts, spectral parameters of the prompt emission, and T_a . By using T_a as a jet angle indicator, and then estimating the collimation corrected prompt energetics, we find a correlation between the latter quantity and the peak energy of the prompt emission. However, this correlation has a large dispersion, similar to the dispersion of the Amati correlation and it is not parallel to the Ghirlanda correlation. Furthermore, we show that the correlation itself results mainly from the dependence of the jet opening angle on the isotropic prompt energy, with the time T_a playing no role, contrary to what we find for the jet break time t_{jet} . We also find that for the bursts in our sample T_a weakly correlates with $E_{\gamma,\text{iso}}$ of the prompt emission, but that this correlation disappears when considering all bursts of known redshift and T_a . There is no correlation between T_a and the isotropic energy of the plateau phase.

Key words: gamma rays: bursts, X–rays: general, radiation mechanisms: general.

1 INTRODUCTION

Before *Swift*, the afterglow observations were typically starting only several hours after the burst, when the flux typically showed a single power law decay. The detection of GRB afterglows provided strong confirmation for the fireball shock model (Rees & Meszaros 1992), which explains it in terms of forward shock emission running into the circumburst medium. In many of the well sampled afterglows, the steepening in the light curve at late times ($\sim 10^5$ s) has been attributed to a narrow conical jet of semiaperture opening angle θ_j whose edges become visible as it decelerates and widens (e.g. Rhoads et al. 1997). The deceleration is due to the expansion of the GRB outflow into the interstellar medium (ISM) whose properties are still poorly understood. By assuming a uniform density ISM the widely dispersed isotropic energy $E_{\gamma,\text{iso}}$ of GRBs considerably clusters if corrected for the collimation factor $f = (1 - \cos \theta_j)$ (Frail et al. 2001). Later, Ghirlanda, Ghisellini & Lazzati (2004; hereafter GGL04) discovered that the collimation corrected GRB energies $E_\gamma = E_{\gamma,\text{iso}}f$ are tightly correlated with the rest frame peak energy E_p of the νF_ν prompt spectrum (so called “Ghirlanda” correlation). Nava et al. (2006; hereafter N06) discovered that, in the case of a wind–like circumburst density pro-

file (which should be more likely if the progenitor is a massive star), this correlation still exists and becomes linear. Liang & Zhang (2005) discovered the phenomenological counterpart of these correlations showing that a tight correlation exists between the $E_{\gamma,\text{iso}}$, E_p and t_{jet} (observed in the optical light curves) computed in the source frame. N06, indeed, demonstrated that this phenomenological correlation is consistent with the E_p – E_γ correlations computed either in the case of a homogeneous and wind medium (HM and WM, respectively).

The jetted–fireball model predicts that when the bulk Lorentz factor $\Gamma \sim 1/\theta_j$, due to the deceleration of the expanding fireball, an achromatic break appears in the afterglow light curve. Such achromaticity is what distinguishes a jet break (i.e. due to the geometry of the event) from a spectral break due to the time–evolution of the characteristic frequencies of the spectrum. Most of the jet breaks of the pre–*Swift* GRBs were, indeed, achromatic, but in a narrow frequency band, from the near–IR to the optical. Moreover, the typical jet breaks were observed between several hours to few days after the GRB trigger: in order to identify this transition in the optical light curves a systematic follow up of the optical afterglow, up to several days after the jet break, was required.

It was highly expected that the multi–wavelength *Swift* satellite (Gehrels et al. 2004) would clearly detect achromatic X–ray to optical breaks. Instead, *Swift* discovered that the X–ray light curve

* E–mail: lara.nava@brera.inaf.it

of GRBs is much more complex than thought (e.g. Burrows et al. 2005). The “canonical” X-ray light curve disclosed by the *Swift* observations is composed by an initial steep decay phase (commonly interpreted as the prompt emission tail due to off axis radiation of a switching-off fireball) followed by a shallow decay phase. At the end of this second phase there is a break that marks the beginning of the “normal” decay phase, observed also before *Swift* (i.e. when the X-ray follow up – mostly with *BeppoSAX* – started, several hours after the trigger). In addition, *Swift* found that the X-ray afterglow is characterised by several early and late times flares superposed to the typical power law flux decay (Nousek et al. 2006, O’Brien et al. 2006). In several *Swift* bursts the optical light curve does not track the X-ray one while in some cases flares and multiple breaks have been observed in the optical too. Due to the richness and complexity of the light curves, the identification of a break as a jet break requires care (Ghirlanda et al. 2007 - G07 hereafter).

Some interpretations of the complex multi-break X-ray light curve of *Swift* GRBs have been proposed: while it is commonly accepted that the early steep decay can be interpreted as due to the curvature effect of a switching-off fireball (see e.g. Kumar & Panaitescu 2000), the origin of the intermediate shallow decay is still controversial. It has been proposed (Ghisellini et al. 2007) that flat X-ray could be “late prompt” emission due to a late central engine activity producing shells with a time-decaying bulk Lorentz factor Γ . In this scenario the flat-to-steep transition occurs when Γ of these late shells becomes $\sim 1/\theta_j$.

Recently, Willingale et al. (2007; hereafter W07) interpreted the steep-flat-steep X-ray light curves as due to the superposition of two separate and independent components: the prompt emission (interpreted as internal dissipation) produces the first steep decay while the afterglow emission (external dissipation) is responsible for the shallow and the final steep components. The modelling, introduced by W07, identifies an early break (typically $\sim 10^3$ – 10^4 s) in the X-ray light curve, called T_a , that marks the shallow-to-steep transition. W07 show that if T_a is used (with the same formalism adopted for t_{jet} in the HM case) to compute an angle $\theta_{j,a}$, the collimation corrected energy $E_{\gamma,a} = E_{\gamma,iso}(1 - \cos \theta_{j,a})$ is correlated with E_p . W07 argue that this correlation has the same slope of the $E_p - E_\gamma$ (Ghirlanda) correlation (found with the jet break time t_{jet}). The two correlations are simply displaced by a factor ~ 26 in E_γ . This implies that the two break times T_a and t_{jet} are proportional (i.e. $t_{jet} \sim 90T_a$). In the sample of W07 only GRB 050820A has both T_a and t_{jet} measured, and indeed in this case $t_{jet} \sim 100T_a$. W07 conclude that if there is a link between T_a and t_{jet} , it is doubtful that both times are actually jet break times. However, W07 stress that it is possible that the end of the plateau phase (corresponding to T_a) depends on the total energy of the outflow. They also point out that it would be interesting to study the phenomenological counterpart of the $E_p - E_{\gamma,a}$ correlation similarly to what has been done with the Liang-Zhang correlation with respect to the Ghirlanda one.

The scope of this paper is to investigate the role of the break time T_a in defining the $E_p - E_{\gamma,a}$ correlation and if a Liang-Zhang like correlation exists between the quantities E_p , $E_{\gamma,iso}$ and T'_a (where E_p and T'_a are expressed in the source rest frame). To this aim, we enlarge the sample used by W07 from 14 to 23 GRBs and derive the collimated corrected energy $E_{\gamma,a}$ using T_a as a jet break and we reproduce the correlation between E_p and $E_{\gamma,a}$ and test, through the scatter of these correlations, the nature of T_a .

We adopt a standard cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = h = 0.7$.

2 THE SAMPLE

In order to build the $E_p - E_{\gamma,a}$ correlation we need the redshift z , the peak energy of the prompt emission spectrum E_p and, from the X-ray light curve, the time T_a . We found, up to November 2006, 16 objects with all these quantities published in the literature. We added 7 more GRBs (050126 050505 050908 060223A 060522 060607A 060908) for which we analysed the *Swift*-BAT (15–150 keV) spectrum and could satisfactorily constrain the spectral peak energy E_p fitting the spectrum with a cut-off power law model. In these cases we also tested our results (by the F -test) finding that the fit with a cut-off power law is better than a single power law. The details of the fitting, together with the results for other *Swift* bursts, will be given elsewhere (Cabrera et al. 2007 in preparation).

The sample of W07 which is used to build the $E_p - E_{\gamma,a}$ correlation contains 14 long GRBs. However, it is hard to identify those bursts present in both samples because W07 do not give the names of those bursts they use to compute the $E_p - E_{\gamma,a}$ correlation.

Our sample is composed by 23 GRBs whose properties are listed in Table 1. In most cases the value of T_a is taken from the compilation of W07, except for two cases:

- for GRB 050922C we have found that the X-ray light curve does not clearly show the shallow part and, therefore, the value of T_a is hardly constrained. A still acceptable fit is obtained by setting $T_a \sim 10^4$ seconds (a factor 26 larger than what derived by W07 – see G07);
- for GRB 060124 the fit of the X-ray light curve should account for the late time achromatic break (Romano et al. 2006; Curran et al. 2006) which corresponds to the jet break time (G07). With the inclusion of this further late time break $T_a \sim 5000$ s (a factor 8 smaller than what reported in W07). The fit is shown in Fig. 1.

Moreover, since the sample of W07 comprises GRBs up to August 2006, for GRB 060908, GRB 060927 and GRB 061121, we have estimated T_a following the procedure outlined in W07, and using the publicly available X-ray light-curves¹. The corresponding light curves and fitting models are shown in Fig. 1.

In the following sections we analyse the $E_p - E_{\gamma,a}$ correlation (found using T_a as an angle estimator) and compare it with the $E_p - E_\gamma$ correlation (found using t_{jet} as angle indicator). We also study the relevance of either T_a and t_{jet} in “collapsing” the scatter of the Amati correlation (Amati et al., 2002; Amati 2006), in the respective collimation corrected correlations.

For the $E_p - E_{\gamma,a}$ correlation we use the sample of 23 GRBs of the *Swift*-era reported in Tab. 1. For the comparison with the $E_p - E_\gamma$ correlation we use the most updated sample of 23 bursts for which t_{jet} has been firmly identified from the optical light curve reported in G07². Within this sample (with t_{jet}) there are 6 GRBs of the *Swift*-era for which we also have an estimate of T_a . To reproduce the Amati correlation (which only requires the knowledge of the redshift and of the spectral parameters) we have used those GRBs that appear at least in one of the two samples. This larger sample contains 40 GRBs.

¹ <http://astro.berkeley.edu/~nat/swift/> by N. Butler, see also Butler & Kocevski (2007)

² In the present analysis we do not include the lower limits on t_{jet} discussed in G07.

GRB	z	E_p keV	$E_{\gamma, \text{iso}, 52}$ erg	Instr ^a	Ref ^b	$\log T_a^c$ s	Ref ^d
050318	1.44	115±27	2.03±0.34	SWI	1	2.01 [1.37–2.68]	W07
050401	2.90	501±117	41±8	KON	2 ^e	3.87 [3.69–4.11]	W07
050416A	0.653	28.6±8.3	0.083±0.029	SWI	3	3.19 [2.69–3.48]	W07
050525A	0.606	127±5.5	2.96±0.64	SWI	4	2.92 [1.69–3.06]	W07
050603	2.821	1333±107	59.8±4.0	KON	5	4.83 [3.69–5.13]	W07
050820A	2.612	1325±277	97.8±8.0	KON	6	3.96 [3.84–4.04]	W07
050922C	2.198	417±118	4.61±0.87	HET	7	2.58 [2.48–2.69]	W07
...	...	...	...	...	...	3.98 [3.68–4.16]	G07
051109A	2.346	539±381	7.58±0.94	KON	8	3.93 [3.70–4.08]	W07
060115	3.53	281±82	8.0±1.5	SWI	9	3.86 [2.86–5.47]	W07
060124	2.297	636±162	43±4.0	KON+SWI	10	4.60 [4.47–4.69]	W07
...	...	...	...	...	...	3.70 [3.40–3.88]	this paper, see Fig. 1
060206	4.048	381±98	4.75±0.76	SWI	11	3.86 [3.68–4.00]	W07
060418	1.489	572±114	12.8±1.1	KON	12	3.44 [3.26–3.58]	W07
060526	3.21	105±21	2.58±0.26	SWI	13	3.84 [3.51–4.31]	W07
060707	3.43	292±71	6.83±1.49	SWI	14	3.58 [2.87–4.05]	W07
060927	5.6	473±116	9.69±1.62	SWI	15	3.60 [3.30–3.78]	this paper, see Fig. 1
061121	1.314	1289±153	26.1±3.0	KON/RHE	16–17	3.90 [3.60–4.08]	this paper, see Fig. 1
050126	1.29	202±49	1.10±0.30	SWI	this paper	2.34 [1.34–5.64]	W07
050505	4.27	622±211	19.5±3.1	SWI	this paper	4.39 [4.15–4.87]	W07
050908	3.344	191±40	2.05±0.34	SWI	this paper	3.31 [2.03–4.17]	W07
060223A	4.41	341±69	4.54±0.72	SWI	this paper	2.73 [2.37–3.05]	W07
060522	5.11	415±78	8.02±1.65	SWI	this paper	2.86 [2.61–3.32]	W07
060607A	3.082	535±164	12.2±1.8	SWI	this paper	4.75 [4.73–4.78]	W07
060908	2.43	487±116	9.17±1.57	SWI	this paper	3.10 [2.80–3.28]	this paper, see Fig. 1

Table 1. GRB sample used to study the $E_p - E_{\gamma, \text{a}}$ correlation. The peak energy E_p is in the source rest frame. The time T_a is in the observer frame. $E_{\gamma, \text{iso}}$ is given in units of 10^{52} erg. ^aThe + sign indicates that the spectral parameters are derived from the joint spectral fit of two different instruments. A sign / indicates that the reported values of E_p and $E_{\gamma, \text{iso}}$ are an average of the values found by two different instruments. The instruments are: SWI=Swift/BAT, KON=Konus/Wind, HET=HETE/Fregate, RHE=RHESSI. ^bReferences for the spectral parameters: (1) Perri et al., 2005; (2) Golenetskii et al., 2005a; (3) Sato et al., 2006; (4) Blustin et al., 2006; (5) Golenetskii et al., 2005b; (6) Cenko et al., 2006; (7) Crew et al., 2005; (8) Golenetskii et al., 2005c; (9) Barbier et al., 2006; (10) Romano et al., 2006; (11) Palmer et al., 2006; (12) Golenetskii et al., 2006a; (13) Schaefer 2007; (14) Stamatikos et al., 2006a; (15) Stamatikos et al., 2006b; (16) Golenetskii et al., 2006b; (17) Bellm et al., 2006. ^cValues in square brackets are the lower and the upper 90% confidence limits. ^dReferences for T_a : Willingale et al. 2007 (W07); Ghirlanda et al. 2007 (G07). ^eThe values of E_p and $E_{\gamma, \text{iso}}$ derive from the average of the two spectral fit presented in the corresponding reference. For GRB 050922C and 060124 we report the value of T_a given in W07 (first line) and the value obtained in G07 and this paper (second line), respectively.

3 ANALYSIS AND RESULTS

First we study the Ghirlanda like correlation and the Liang & Zhang like correlation defined using T_a in place of t_{jet} . We use the formalism defined in N06 where, for a generic time break t_{break} , the jet opening angle is defined as³:

$$\begin{aligned} \theta &= 0.161 \left(\frac{t_{\text{break}}}{1+z} \right)^{3/8} \left(\frac{n \eta_\gamma}{E_{\gamma, \text{iso}, 52}} \right)^{1/8} \quad [\text{HM}] \\ \theta &= 0.2016 \left(\frac{t_{\text{break}}}{1+z} \right)^{1/4} \left(\frac{\eta_\gamma A_*}{E_{\gamma, \text{iso}, 52}} \right)^{1/4} \quad [\text{WM}] \end{aligned} \quad (1)$$

where n represents the circumburst density in the homogeneous case (HM) and A_* is the normalisation of the wind (WM) density profile ($n(r) \propto A_* r^{-2}$ – see N06 for more details). Here η_γ is the radiative efficiency of the prompt phase assumed to be 0.2 for all bursts.

In both cases of Eq. 1, it is necessary to know the isotropic bolometric rest frame equivalent energy $E_{\gamma, \text{iso}}$. This is

$$E_{\gamma, \text{iso}} = \frac{4\pi d_L^2 S_\gamma k}{1+z} \quad (2)$$

³ We use the notation $Q_x = 10^x Q$, using cgs units.

where d_L is the luminosity distance, S_γ is the γ -ray fluence in the observed energy band, k is the bolometric correction factor needed to find the energy emitted in a fixed energy range (here, 1–10⁴ keV) in the rest frame of the source. For most pre-Swift bursts the spectrum is fitted with the Band function (Band et al. 1993) composed by two smoothly joint power laws. On the other hand, several Swift bursts, due to the narrow energy range of the BAT instrument (15–150 keV), have a spectrum best fitted with a cutoff-power law model. As discussed in Firmani et al. (2006), in these cases the extrapolation of the cutoff-power law model up to 10⁴ keV may considerably underestimate the energy, if the spectrum is, instead, a Band function. To account for this limitation of the spectral fits in the Swift era, the value of $E_{\gamma, \text{iso}}$ reported in Tab. 1 is an average between the value derived with the cutoff-power law model and that derived assuming a Band model with high energy photon index $\beta = -2.3$ (see also G07).

3.1 Correlation analysis

For the 23 GRBs of Tab. 1 we computed the angle $\theta_{j, \text{a}}$, where the subscript “a” means that the angle has been estimated using T_a as a jet break time, and the collimation corrected energy $E_{\gamma, \text{a}}$ in both the WM and HM case. The density in the homogeneous case is




Figure 1. The early BAT and late XRT light curves of GRB 060124, GRB 060908, GRB 060927 and GRB 061121 modelled with the function proposed by W07. The two model components are shown by the long-dashed lines while their sum is shown by the solid line. The fits were performed excluding the flares of the light curves. The long dashed vertical lines show the T_a in each light curve (for GRB 060124 we also report the jet break time found in the optical light curve – see text).

known only for some GRBs and for the others we take the value $n = 3 \pm 2.76$; the density in the wind case is assumed equal for all GRBs and without error (see N06 and G07 for details).

We report also the correlation defined with the jet break time t_{jet} and recently updated with the *Swift* bursts in G07. Here we use the sample of G07 (Tab. 1 in that paper) excluding bursts with only a lower limit on t_{jet} . For comparison we also recomputed the $E_p - E_{\gamma, \text{iso}}$ (Amati) correlation with the sample of 40 GRBs.

The fit is performed using the routine *fitxxy* (Press et al. 1999) which weights for the errors on the involved variables. All the fit results can be expressed in the synthetic form:

$$\left(\frac{E_p}{100 \text{ keV}} \right) = (K \pm \sigma_K) \left(\frac{E_{\gamma, a}}{E_{\text{bar}}} \right)^{s \pm \sigma_s} \quad (3)$$

where $E_{\gamma, a}$ must be replaced by E_γ or $E_{\gamma, \text{iso}}$ according to the correlation considered. Here E_{bar} represents the barycentre of the energy values of the sample considered.

The three correlations are shown in Fig. 2 (homogeneous case) and in Fig. 3 (wind case), and the fit results are listed in Tab. 2. For every correlation we report also the reduced χ^2 and the 1σ scatter of the data points around the best fit line, modelled with a gaussian distribution.

From the comparison of the results reported in Tab. 2 and in Figs. 2 and 3 we note:

(i) the correlation $E_p - E_{\gamma, a}$ derived using T_a as a jet break time is not parallel to the $E_p - E_\gamma$ correlation (derived using t_{jet}). In particular, independently from the specific density profile considered, the $E_p - E_{\gamma, a}$ correlation is much flatter than the $E_p - E_\gamma$ correlation and in the homogeneous case it is even slightly flatter than the Amati correlation. By reconstructing the sample of W07 (through the quoted GCNs and the values given in that paper), and analyzing this smaller sample, we indeed obtain a $E_p - E_{\gamma, a}$ correlation parallel to the $E_p - E_\gamma$ one, although not as tight. It is the inclusion of the additional GRBs that changes the slope of $E_p - E_{\gamma, a}$ correlation and further increases its scatter. Also the 6 GRBs with both T_a and t_{jet} (encircled symbols) seem to define a $E_p - E_{\gamma, a}$ parallel to the $E_p - E_\gamma$ correlation, and this issue will be discussed in Sec. 3.2.

(ii) the fit of the $E_p - E_{\gamma, a}$ correlation (in terms of χ^2) is significantly worse than that of the $E_p - E_\gamma$ correlation, which, instead, is statistically acceptable;

(iii) the $E_p - E_{\gamma, a}$ correlation has a scatter which is a factor 2 larger than that of the $E_p - E_\gamma$ correlation and is similar to that of the Amati correlation.

Item (i) suggests that the relation between t_{jet} and T_a , if any, is not trivial (i.e. t_{jet} is not simply proportional to T_a as suggested by W07). The large scatter of the $E_p - E_{\gamma, a}$ correlation, which is comparable to that of the $E_p - E_{\gamma, \text{iso}}$ correlation, raises the ques-




Figure 2. Homogeneous circumburst medium. From right to left we show the $E_p - E_{\gamma, \text{iso}}$ correlation (filled squares – 40 GRBs), the $E_p - E_{\gamma}$ correlation (filled circles – 23 bursts from G07) and the $E_p - E_{\gamma, \text{a}}$ correlation (filled stars – 23 GRBs of Tab. 1). The shaded regions (for the three correlations) represent the 1, 2, 3 σ scatter of the data points (computed perpendicular) with respect to the best fitted correlation (solid lines). Circled points (on the three correlations) represent the 6 *Swift* bursts for which both T_{a} and t_{jet} are known.




Figure 3. Wind circumburst medium. Symbols as in Fig.2

Correlation	Density profile	N	K (σ_K)	s (σ_s)	E_{bar} (erg)	χ_r^2	Scatter
$E_p - E_{\gamma,a}$	homog.	23	4.71 (0.37)	0.55 (0.05)	3.11×10^{49}	1.95 (21 d.o.f.)	0.16
$E_p - E_{\gamma,a}$	wind	23	4.95 (0.38)	0.78 (0.07)	4.14×10^{49}	1.95 (21 d.o.f.)	0.16
$E_p - E_{\gamma}$	homog.	23	2.89 (0.16)	0.71 (0.04)	4.14×10^{50}	1.14 (21 d.o.f.)	0.09
$E_p - E_{\gamma}$	wind	23	3.21 (0.17)	1.06 (0.07)	2.47×10^{50}	0.91 (21 d.o.f.)	0.08
$E_p - E_{\gamma,\text{iso}}$	—	40	3.77 (0.11)	0.62 (0.02)	1.31×10^{53}	5.70 (38 d.o.f.)	0.16
$E_p - E_{\gamma,\text{iso}}$	—	23	4.83 (0.21)	0.65 (0.03)	1.22×10^{53}	2.89 (21 d.o.f.)	0.15

Table 2. Best fit results of the correlations involving the rest frame peak energy and the collimation corrected energy computed either using T_a as a jet break time and the real jet break time t_{jet} . We also report the Amati correlation for the considered 40 GRBs and for the 23 defining also the $E_p - E_{\gamma,a}$ correlation. The collimation corrected correlations are reported for both the homogeneous and wind medium scenario. K and s represent the normalisation and slope, respectively, of the best fit correlation (see Eq. 3). The 1σ scatter of the data points around the best fit correlation is also reported.

tion whether the new added variable T_a can be treated as a jet break time or not. If not, we are still left with the possibility that a completely phenomenological correlation between E_p and $E_{\gamma,\text{iso}}$ and T'_a exists (where T'_a is the time in the source rest frame). This possibility will be explored in the next section.

3.2 The $E_p - E_{\gamma,\text{iso}} - T'_a$ correlation

In this section we study the $E_p - E_{\gamma,\text{iso}} - T'_a$ correlation. First we fit the multi-variable correlation with the 23 GRBs of Tab. 1 through a least-square method without considering the errors on the three variables. The result is:

$$E_{\gamma,\text{iso},52} = (7.29 \pm 0.89) \left(\frac{E_p}{357 \text{ keV}} \right)^{1.5 \pm 0.1} \times \left(\frac{T'_a}{0.012 \text{ s}} \right)^{0.06 \pm 0.59}. \quad (4)$$

Note that the exponent of T'_a is very close to zero and this implies that T'_a does not reduce the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation. Indeed, the above result is almost identical to the $E_p - E_{\gamma,\text{iso}}$ correlation defined by the 23 GRBs reported in Tab. 1 (see also the last row of Tab. 2).

For consistency with the fits reported in the previous sections we also fitted the $E_p - E_{\gamma,\text{iso}} - T'_a$ correlation by weighting for the errors on the three variables. In this case we find a different result:

$$E_{\gamma,\text{iso},52} = (13.8 \pm 2.0) \left(\frac{E_p}{513 \text{ keV}} \right)^{2.18 \pm 0.53} \times \left(\frac{T'_a}{0.0173 \text{ s}} \right)^{-0.67 \pm 0.43}. \quad (5)$$

with a reduced $\chi^2 = 2.28$ (20 dof). The reason for the difference with Eq. 4 lies in the errors on T'_a , which are particularly large (see Tab. 1). To prove it, we have reduced the errors on T'_a by a factor 1.5 and recovered the result of the fit with a simple linear regression analysis (Eq. 4).

In Fig. 4 we compare the scatter of the data points around the $E_p - E_{\gamma,\text{iso}}$ correlation (defined by the 23 GRBs of Tab. 1) with the scatter of the data points around the best fit plane of Eq. 5 (i.e. obtained by weighing for the errors). We note that the scatter of the data points in the $E_p - E_{\gamma,\text{iso}} - T'_a$ plane is comparable to the scatter of the data points in the $E_p - E_{\gamma,\text{iso}}$ plane. This suggests that, although in Eq. 5 (found by weighting for the errors) the exponent of T'_a is different from zero, the scatter of the data points is not improved with respect to the $E_p - E_{\gamma,\text{iso}}$ correlation. We conclude that it is Eq. 4 which better represents the best fit to the data.

This would be sufficient to demonstrate that T_a does not help




Figure 4. Scatter distribution of the 23 GRBs of Tab. 1 around the best fit $E_p - E_{\gamma,\text{iso}}$ correlation (top histogram) compared with the scatter of the same points around the best fit plane of the $E_p - E_{\gamma,\text{iso}} - T'_a$ correlation defined by Eq. 5 (bottom histogram). One can see that the scatter of the $E_p - E_{\gamma,\text{iso}} - T'_a$ correlation is similar (if not larger) than the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation.

to define a tighter correlation between the isotropic energy and E_p . To further demonstrate this point, we directly study the relation between T_a and $E_{\gamma,\text{iso}}$, comparing it with the corresponding relation between t_{jet} and $E_{\gamma,\text{iso}}$.

Consider bursts with the same E_p . For these the existence of the empirical Liang & Zhang correlation implies that $E_{\gamma,\text{iso}} \propto t'_{\text{jet}}{}^{-1}$. This is proved by the data: in Fig. 5 we show the anticorrelation between t_{jet} (rest frame) and $E_{\gamma,\text{iso}}$ for the sample of 23 GRB by considering bursts with similar E_p (corresponding to the different symbols in Fig. 5). The $E_{\gamma,\text{iso}} \propto t'_{\text{jet}}{}^{-1}$ relation converts in $E_{\gamma,\text{iso}} \theta_j^2 = \text{const}$ (the “Frail correlation”) when using the relation between t_{jet} and θ_j (Eq. 1), for both the HM and the WM cases.

Consider now Fig. 6, showing T'_a versus $E_{\gamma,\text{iso}}$: in this case T'_a does not anticorrelate with $E_{\gamma,\text{iso}}$, for constant E_p . The absence of the anticorrelation translates in the impossibility to obtain a similar “Frail correlation” when using T_a as a jet angle indicator (i.e. inserting T_a instead of t_{jet} in Eq. 1).




Figure 5. The time t_{jet} as a function of $E_{\gamma,\text{iso}}$ for the burst of the G07 sample. Bursts with different peak energy ranges (as labelled) are shown with different symbols. The dotted lines indicate $t_{\text{jet}} \propto E_{\gamma,\text{iso}}^{-1}$, with different normalizations, to illustrate that bursts with the same E_p do follow this law.




Figure 6. The time T'_a vs $E_{\gamma,\text{iso}}$ for the burst of this paper (Tab. 1). As in Fig. 5 we use different symbols for bursts with E_p in different ranges.

3.3 The origin of the $E_p - E_{\gamma,a}$ correlation

Although T_a spans 3 orders of magnitudes, and although it is not a jet angle indicator, the $E_p - E_{\gamma,a}$ correlation exists and has a scatter comparable (or slightly larger) than the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation. What is, then, its origin?

In Eq. 1 the jet angle depends on both the break time (either T_a or t_{jet}) but also on $E_{\gamma,\text{iso}}/\eta_\gamma$. This term is the kinetic energy of the GRB outflow left-over after the prompt radiative phase. Since, usually, one assumes a constant value $\eta_\gamma = 0.2$, the collimation correction depends on $E_{\gamma,\text{iso}}$.

In this section we explore the role of $E_{\gamma,\text{iso}}$ and t_{break} (either identified as T_a or t_{jet}) in Eq. 1 in shaping the $E_p - E_{\gamma,a}$ and the $E_p - E_\gamma$ correlation and their scatter.

From Eq. 1 considering only the dependencies from $E_{\gamma,\text{iso}}$

and t_{break} we find (under the assumption of small angles):

$$\begin{aligned} \text{WM} : E_\gamma &\propto E_{\gamma,\text{iso}} \theta^2 \propto E_{\gamma,\text{iso}} \frac{t_{\text{break}}^{1/2}}{E_{\gamma,\text{iso}}^{1/2}} \propto E_{\gamma,\text{iso}}^{1/2} t_{\text{break}}^{1/2} \\ \text{HM} : E_\gamma &\propto E_{\gamma,\text{iso}} \theta^2 \propto E_{\gamma,\text{iso}} \frac{t_{\text{break}}^{3/4}}{E_{\gamma,\text{iso}}^{1/4}} \propto E_{\gamma,\text{iso}}^{3/4} t_{\text{break}}^{3/4} \end{aligned} \quad (6)$$

Consider those GRBs that have the same E_p , but different $E_{\gamma,\text{iso}}$. Even if they have the same t_{break} , the use of Eq. 6 implies a clustering of the corresponding collimation corrected E_γ : GRB with larger $E_{\gamma,\text{iso}}$ would have smaller θ , and therefore have a larger correction for their collimation (and vice-versa).

We calculate θ and correct for the collimation factor assuming that the break time (either T_a or t_{jet}) is constant and equal for all bursts. The use of Eq. 6 with a fixed t_{break} gives $E_\gamma \propto E_{\gamma,\text{iso}}^{1/2}$ in the WM case, and $E_\gamma \propto E_{\gamma,\text{iso}}^{3/4}$ in the HM case. We plot E_p versus this estimate of E_γ in Fig. 7 (for both the WM and HM cases). The empty star symbols define a correlation which is less scattered than the $E_p - E_{\gamma,\text{iso}}$ correlation (empty circles) in the HM and WM case (left and right panels in Fig. 7). Moreover, the scatter reduction is somewhat more significant in the WM case with respect to the HM case.

Note that this result has no physical meaning and it is only due to the fact that we plot the points in the $E_p - E_{\gamma,\text{iso}}^{1/2}$ plane for the WM case (or $E_p - E_{\gamma,\text{iso}}^{3/4}$ for the HM case) instead of in the $E_p - E_{\gamma,\text{iso}}$ plane. Up to this point the reduction of the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation, and the difference in the slope with respect to the Amati relation, is due to the dependence of Eq. 6 (used to derive the jet opening angle) on $E_{\gamma,\text{iso}}$.

We can now see if the use of the real break time (either T_a or t_{jet} – upper panels and lower panels in Fig. 7, respectively) improves or not the found correlations. The results in the case of T_a are shown in the upper panels of Fig. 7 for the HM and WM case. Although a correlation between E_p and $E_{\gamma,a}$ appears (filled circles), we find that the effect of T_a is to increase the scatter of the data points with respect to the same correlation defined with T_a constant. Note that the scatter of the found $E_p - E_{\gamma,a}$ correlation is not less than that of the $E_p - E_{\gamma,\text{iso}}$ correlation (both in the HM and WM case). This demonstrates that the variable T_a , if treated as a jet break time to estimate the angle and derive the collimation corrected energy, leaves unaltered (or even increases) the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation. This effect is clearly shown in Fig. 7 by bursts with typical rest frame peak energy in the range 100–1000 keV which are the majority in our sample.

In the bottom panels of Fig. 7 we show the results obtained with t_{jet} . As discussed in Sec. 2, the sample of bursts used to define the $E_p - E_\gamma$ correlation only marginally overlaps to that used to define the $E_p - E_{\gamma,a}$ correlation: this is the reason why also the $E_p - E_{\gamma,\text{iso}}$ correlation in the lower and upper panels of Fig. 7 are different. If first we set $t_{\text{jet}} = \text{constant}$ and equal for all bursts we have, as before, a reduction (empty stars) of the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation (empty circles). But now, if we assign to each GRB in our sample its t_{jet} found from the optical light curve, we see that the scatter of the data points is further reduced (note in particular this effect in peak energy range 500–1000 keV).

From these results we conclude that:

- the dependence of the jet angle on $E_{\gamma,\text{iso}}$ reduces the scatter of the $E_p - E_{\gamma,\text{iso}}$ correlation;
- the use of the time T_a which marks the end of the plateau phase in the X-ray light curves, increases the scatter of the correla-




Figure 7. Comparison of the scatter of the $E_p - E_{\gamma,a}$ and $E_p - E_\gamma$ correlations in the homogeneous (left) and wind (right) medium cases. Top panels: from right to left we report the $E_p - E_{\gamma,iso}$ correlation (empty circles), the correlation defined by assuming a constant value of T_a (for all the bursts) in computing the collimation correction (empty stars) and the correlation defined by using the measured T_a (filled circles; values of T_a reported in Tab. 1). Bottom panels: same as above but considering t_{jet} instead of T_a .

tion found assuming a constant T_a . This effect, counterbalanced by the previous one, makes the scatter of the $E_p - E_{\gamma,a}$ correlation similar to the scatter of the $E_p - E_{\gamma,iso}$ correlation;

- the (albeit weak) positive correlation between T_a and $E_{\gamma,iso}$ (see Fig. 6), makes the slope of the $E_p - E_{\gamma,a}$ correlation to differ somewhat from the slope of the correlation found with T_a constant;
- the small dispersion of the Ghirlanda relation is due not only to the presence of $E_{\gamma,iso}$ in the equation for the angle, but also to the jet break time t_{jet} .

4 SUMMARY AND DISCUSSION

We have analyzed in detail the correlation found by W07 between E_p and the collimation corrected energy $E_{\gamma,a}$, found using T_a as a jet break. Our sample (of 23 objects) is larger than what used by W07, because we could add some GRB occurred after August 2006, and some other bursts for which we could find the value of E_p . We find that E_p and $E_{\gamma,a}$ correlate, but we also find that this

correlation is weaker, and has a different slope, than the $E_p - E_\gamma$ one.

Investigating further, we have also shown that the $E_p - E_{\gamma,a}$ correlation is entirely a consequence of the existence of the $E_p - E_{\gamma,iso}$ (Amati) correlation, coupled with the fact that the opening angle θ_j depends on $E_{\gamma,iso}$. This is why one finds a correlation which is narrower than the Amati one even using the same T_a for all bursts: *in this case the found correlation is even better than the one obtained introducing the real T_a* . This demonstrates that T_a does not play any role in the $E_p - E_{\gamma,a}$ correlation: it is even worsening it.

We have then made the very same test to the $E_p - E_\gamma$ (Ghirlanda) correlation, and found that in this case the use of the “real” t_{jet} improves the correlation, decreases the scatter and slightly changes the slope of the correlation, with respect to the one obtained using a fixed t_{jet} .

All these results are fully confirmed by analyzing the $E_p - E_{\gamma,iso} - T'_a$ correlation, and comparing it to the Liang & Zhang correlation (which uses t_{jet} instead of T_a). Again, all evidence is to-




Figure 8. Top panel: the time T'_a as a function of the isotropic energy of the prompt emission $E_{\gamma,iso}$ calculated in the 15–150 keV energy range, without including any bolometric and k -correction for all bursts with measured redshift (circles) and for those which also have a measured E_p (stars). The insert shows the bolometric fluence vs the 15–150 keV fluence, to show that the two are proportional. Bursts in the insert are the ones in Tab. 1. Bottom panel: the time T'_a as a function of the energy emitted in the plateau phase, in the 0.3–10 keV energy range. The fluences of the plateau phase have been taken from W07. Symbols as above.

wards T'_a playing no role: the found $E_p - E_{\gamma,iso} - T'_a$ correlation has a similar (if not larger) scatter than the $E_p - E_{\gamma,iso}$ correlation.

We conclude that the time T_a at which the plateau of the X-ray light curve ends is not an important parameter for defining the spectral–energy correlations. As a consequence, it is not a jet angle indicator. Instead, we confirm that t_{jet} is likely to measure the jet opening angle, since it passed our simple test to see if it helps or not to strengthen the correlation between E_p and the collimation corrected energy derived using a fixed value of t_{jet} for all bursts.

We note a weak correlation between T'_a and $E_{\gamma,iso}$ when considering the 23 GRBs in our sample (see Fig. 6, probability of chance correlation equal to 4×10^{-3} , according to the Spearman rank correlation), which however disappears when considering the entire sample of GRBs of known redshifts and T'_a (and relaxing the requirement to have also E_p known, see Fig. 8). Therefore we can also conclude that T'_a is not related to $E_{\gamma,iso}$, differently from t_{jet} which is instead anti-correlated with $E_{\gamma,iso}$ when considering GRBs of similar E_p . Note that, for a distribution of T'_a not correlated (or anti-correlated) with $E_{\gamma,iso}$, the resulting $E_p - E_{\gamma,a}$ correlation should become parallel to the one obtained fixing T_a to a constant value. We therefore predict that, for larger samples of GRBs with known redshift, T_a and E_p , the $E_p - E_{\gamma,a}$ correlation will have a slope ≈ 0.75 (≈ 1) in the HM (WM) case.

Finally, there is no correlation between T'_a and the isotropic

energy (in the observed 0.3–10 keV energy range) emitted during the plateau phase (see Fig. 8).

Unfortunately, our finding that the $E_p - E_{\gamma,a}$ correlation is only a by-product of the $E_p - E_{\gamma,iso}$ (Amati) correlation is not very helpful in shedding new light on the problem of explaining why the light curve of most GRBs is characterized by the steep–flat–steep behaviour, and also why this is sometimes different by the simultaneous behavior observed in the optical band.

Recently, we (Ghisellini et al. 2007) have proposed that the X-ray light curve is produced by internal dissipation in late shells, where “late” means that the central engine continues to create relativistic shells even days after the trigger. This implies that the central engine is long lived, and that there is a sharp transition between the early phase (determining the duration of the early prompt, i.e. the time T_{90}), and the late phase, emitting much smaller luminosity, but for a longer time. The early phase should be characterized by a large bulk Lorentz factor Γ of the created shells, changing erratically, while the late phase is instead characterized by shells created with Γ decreasing monotonically. This implies that when $\Gamma > 1/\theta_j$, we see only a fraction of the emitting surface, but when $\Gamma \leq 1/\theta_j$ we see the entire emitting surface. This means that there should be a break in the light curve when $\Gamma = 1/\theta_j$. We associate this time to T_a .

According to this idea, the two times T_a and t_{jet} have an obvious link: both are determined by the jet opening angle θ_j . Then, why we do not observe a relation between these two times? We propose the following answer: the time t_{jet} is determined when the fireball made by the merged shells of the early phase, carrying a kinetic energy E_k , is decelerated by the external medium. The dynamics of the process is robust, depending only on the value of the external medium density (and its profile with distance), on E_k , and finally on the conservation of energy and momentum. We have the same $\Gamma(t)$ law for different GRBs, if the external medium has the same profile.

On the contrary, the law characterizing how Γ changes as a function of time during the late prompt phase can be different from burst to burst. As far as we know, there might not be a robust analog of the conservation of energy and momentum dictating the $\Gamma(t)$ behaviour in this case. Indeed, we can have information on this law from the observed slopes of the light curve (see Ghisellini et al. 2007), and under simplifying hypotheses we can already derive that $\Gamma(t)$ is different from one bursts to another. As a consequence, there is no general formula, equal to all bursts, in which T_a can be inserted to derive the jet opening angle.

ACKNOWLEDGEMENTS

We thank F. Tavecchio for useful discussions and the anonymous referee for his/her suggestions. PRIN–INAF is thanked for a 2005 funding grant.

REFERENCES

- Amati, L., Frontera, F., Tavani, M., et al. 2002, *A&A*, 390, 81
- Amati, L., 2006, *MNRAS*, 372, 233
- Band D. L., et al., 1993, *ApJ*, 413, 281
- Barbier, L., Barthelmy, S., Cummings, J., et al. 2006, *GCN*, 4518
- Bellm, E., Bandstra, M., Boggs, S., Wigger, C., Hajdas, W., Smith, D.M., & Hurley, K. 2006, *GCN*, 5838
- Blustin, A.J., Band, D., Barthelmy, S., et al. 2006, *ApJ*, 637, 901
- Burrows et al., 2005, *Sci*, 309, 1833

- Butler, N.R. & Kocevski, D., 2007, subm to ApJ (astro-ph/0612564)
- Cenko, S.B., Kasliwal, M., Harrison, F.A., et al. 2006, ApJ, 652, 490
- Crew, G., Ricker, G., Atteia, J-L., et al. 2005, GCN, 4021
- Curran, P., Kann, D.A., Ferrero, P., Rol, E. & Wijers R.A.M.J., 2006, Il Nuovo Cimento C., in press (astro-ph/0611189)
- Firmani et al., 2006, MNRAS, 370, 185
- Frail D. et al., 2001, ApJ, 562, L55
- Gehrels, N., Chincarini, G., Giommi P., et al., 2004, ApJ, 611, 1005
- Ghirlanda, G., Ghisellini, G. & Lazzati, D., 2004, ApJ, 616, 331 (GGL04)
- Ghirlanda, G., Nava, L., Ghisellini, G. & Firmani, C., 2007, A&A subm (G07)
- Ghisellini, G., Ghirlanda, G., Nava, L. & Firmani, C., 2007, ApJ Lett. subm. (astro-ph/0701430)
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., & Cline, T. 2005a, GCN, 3179
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., & Cline, T. 2005b, GCN, 3518
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., & Cline, T. 2005c, GCN, 4238
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., Ulanov, M., & Cline, T. 2006a, GCN, 4989
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., & Cline, T. 2006b, GCN, 5837
- Liang, E. & Zhang, B., 2005, ApJ, 633, L611
- Kumar, P. & Panaitescu, A., 2000, ApJ, 647, 1213
- Nava, L., Ghisellini, G., Ghirlanda, G., Tavecchio, F. & Firmani, C., 2006, A&A, 450, 471 (N06)
- Nousek J., et al., 2006, ApJ, 642, 389
- O'Brien et al., 2006., ApJ, 647, 1213
- Palmer, D., Barbier, L., Barthelmy, S., et al. 2006, GCN, 4697
- Perri, M., Giommi, P., Capalbi, M., et al. 2005, A&A, 442, L1
- Press, W.H. et al. 1999, *Numerical Recipes in C*, Cambridge University Press, 661
- Rees, M.J. & Meszaros, P., 1992, MNRAS 258, L41
- Rhoads J., 1997, ApJ, 487, L1
- Romano, P., Campana, S., Chincarini, G., et al. 2006, A&A, 456, 917
- Sato, G., Yamazaki, R., Ioka, K., et al. 2006, astro-ph/0611148, accepted for publication in ApJ
- Schaefer, B.E. 2007, ApJ, in press (astro-ph/0612285)
- Stamatikos, M., Barbier, L., Barthelmy, S., et al. 2006a, GCN, 5289
- Stamatikos, M., Barbier, L., Barthelmy, S., et al. 2006b, GCN, 5639
- Willingale, R., et al., 2007, subm to ApJ (W07) (astro-ph/0612031)